

YOU ARE INVITED TO A SPECIAL WEBINAR EVENT



NON-OPIOID
EXPAREL[®]
(bupivacaine liposome injectable suspension)

PACIRA
BIOSCIENCES, INC.

LEVELING UP: Leveraging Regional Analgesic Approaches to Elevate Patient Care

An interactive webinar series highlighting practice-changing pain management protocols presented as a 10-minute case study with a 20-minute live Q&A session



Hosted by:

Gary Schwartz, MD

Director of Acute Pain Management of Maimonides Medical Center in Brooklyn, NY

Case Study: Implementing TAP Blocks to Evolve Low-Opioid Protocols in C-Section

Event details:



Join Dr. Gary Schwartz as he discusses implementing a TAP-based regional analgesia program to enable low- or no-opioid C-sections for women in a large metropolitan medical center. Dr Schwartz will also field live questions from the audience.

Date: Wednesday, June 16, 2021

Time: 7:00 PM ET/4:00 PM PT



RSVP:

Register by scanning the QR code or at URL: bit.ly/PaciraLevelUpJune

Questions? Contact the Medical Innovation and Education team member:

Nicole Adams: (404) 376-7627 or nicole.adams@pacira.com



Indication

EXPAREL® (bupivacaine liposome injectable suspension) is indicated for single-dose infiltration in patients aged 6 years and older to produce postsurgical local analgesia and in adults as an interscalene brachial plexus nerve block to produce postsurgical regional analgesia. Safety and efficacy have not been established in other nerve blocks.

Important Safety Information

EXPAREL is contraindicated in obstetrical paracervical block anesthesia.

Adverse reactions reported in adults with an incidence greater than or equal to 10% following EXPAREL administration via infiltration were nausea, constipation, and vomiting; adverse reactions reported in adults with an incidence greater than or equal to 10% following EXPAREL administration via interscalene brachial plexus nerve block were nausea, pyrexia, and constipation.

Adverse reactions with an incidence greater than or equal to 10% following EXPAREL administration via infiltration in pediatric patients six to less than 17 years of age were nausea, vomiting, constipation, hypotension, anemia, muscle twitching, vision blurred, pruritis, and tachycardia.

If EXPAREL and other non-bupivacaine local anesthetics, including lidocaine, are administered at the same site, there may be an immediate release of bupivacaine from EXPAREL. Therefore, EXPAREL may be administered to the same site 20 minutes after injecting lidocaine.

EXPAREL is not recommended to be used in the following patient populations: patients <6 years old for infiltration, patients younger than 18 years old for interscalene brachial plexus nerve block, and/or pregnant patients.

Because amide-type local anesthetics, such as bupivacaine, are metabolized by the liver, EXPAREL should be used cautiously in patients with hepatic disease.

Warnings and Precautions Specific to EXPAREL

Avoid additional use of local anesthetics within 96 hours following administration of EXPAREL.

EXPAREL is not recommended for the following types or routes of administration: epidural, intrathecal, regional nerve blocks **other than interscalene brachial plexus nerve block**, or intravascular or intra-articular use.

The potential sensory and/or motor loss with EXPAREL is temporary and varies in degree and duration depending on the site of injection and dosage administered and may last for up to 5 days, as seen in clinical trials.

Warnings and Precautions for Bupivacaine-Containing Products

Central Nervous System (CNS) Reactions: There have been reports of adverse neurologic reactions with the use of local anesthetics. These include persistent anesthesia and paresthesia. CNS reactions are characterized by excitation and/or depression.

Cardiovascular System Reactions: Toxic blood concentrations depress cardiac conductivity and excitability which may lead to dysrhythmias, sometimes leading to death.

Allergic Reactions: Allergic-type reactions (eg, anaphylaxis and angioedema) are rare and may occur as a result of hypersensitivity to the local anesthetic or to other formulation ingredients.

Chondrolysis: There have been reports of chondrolysis (mostly in the shoulder joint) following intra-articular infusion of local anesthetics, which is an unapproved use.

Methemoglobinemia: Cases of methemoglobinemia have been reported with local anesthetic use.

Full prescribing information is available at www.EXPAREL.com or call 1-855-793-9727.

NON-OPIOID
EXPAREL®
(bupivacaine liposome injectable suspension)

PACIRA
BIOSCIENCES, INC.